PREVENTING SUICIDE IN SÁPMI

Jon Petter Stoor, PhD, clinical psychologist

Epidemiology and Global Health, Umeå University, Sweden
Centre for Sámi Health Research, Community Medicine, UiT – the Arctic University of Norway




WHO AM I AND WHAT IS THE CONTEXT?



OVERVIEW







+MUSIC VIDEOS



KNOWLEDGECOMPILATION ON PSYCHOSOCIAL HEALTH AMONG SÁMI



- Suicide
- Suicidality
- Mental health




STANDARDISED MORTALITY RATIOS FOR SUICIDE AMONG SÁMI COMPARED TO MAJORITY POPULATIONS IN SWEDEN, NORWAY AND FINLAND

	G-1t	\wedge	D.
	Cohort	IVI VII	F
Northern Sweden (1961 - 2001)	Total cohort	1.17	0.76
	Non-herding	1.05	0.67
	Herding	1.50	1.12
Northern Norway (1970 – 1998)	Total cohort	1.27	1.27
	Finnmark	1.50	1.55
	Troms	0.74	1.00
	Nordland	0.42	3.17
	Core area	1.54	1.31
	Coast	1.24	1.21
	South	0.41	1.51
	1970-1980	1.17	1.14
	1981–1990	1.36	1.92
	1991–1998	1.20	0.81
	Non-herding	1.30	1.34
	Herding	1.06	0.66
Northern Finland (1979-2010)	Total cohort	1.78	1.26
	1979-1987	1.83	(No case)
	1988–1996	1.07	1.93
	1997-2005	2.55	1.2
	2006-2010	2.32	1.2

Young, Revich, & Soininen (2015)


SUICIDE AMONG REINDEER HERDING SÁMI IN SWEDEN (1961–2017) BY SEX AND AGE






Jacobsson, Stoor & Eriksson, 2020

SUICIDE METHODS AMONG REINDEER HERDING SÁMI MEN IN SWEDEN (1961–2017)





Jacobsson, Stoor & Eriksson, 2020
UMEÅ UNIVERSITY

SUICIDES BY SEX AND REGION AMONG REINDER ERDING SÁMI IN SWEDEN, 1961-2017




SUICIDAL EXPRESSIONS AMONG REINDEER HERDING SÁMI IN SWEDEN, AND A SWEDISH REFERENCE POPULATION, BY ETHNICITY AND SEX, MEASURED WITH THE ATTITUDES TOWARDS SUICIDE SCALE (ATTS) IN 2007–2008





DEPRESSION AND ANXIETY AMONG REINDEER HERDING SÁMI IN **SWEDEN (HADS CUT-OFFS)**



■ "Northener" rural women (n= 369) ■ Sámi reinder herding men (n= 165)


■ "Northener" urban men (n= 321) ■ "Northener" rural men (n= 324)

UMEÅ UNIVERSITY

Kaiser, N., Sjolander, P., Edin-Liljegren, A., Jacobsson, L., & Salander Renberg, E. (2010)

PERSONS WITH LOW CONFIDENCE IN PRIMARY CARE, PSYCHIATRY AND SOCIAL SERVICES AMONG REINDEER HERDING SÁMI AND «NORTHENER» SWEDES, BY SEX

Daerga et al. (2012)




UMEÅ UNIVERSITY

A SÁMI PERSPECTIVE?

Stoor, JPA., Kaiser, N., Jacobsson, L., Salander-Renberg, E., & Silviken, A. (2015). "We are like lemmings": making sense of the cultural meaning(s) of suicide among the indigenous Sami in Sweden. *International Journal of Circumpolar Health*, *74*, 27669.



MUSIC VIDEO: ÁHPI – SOFIA JANNOK



MAPPING SUICIDE PREVENTION IN SÁPMI



1. Identify suicide prevention initiatives targeting Sámi

- Utilise prior knowledge and networks
- · Acquire project reports and other written documentation

2. Describe initiatives

- Create a template
- Input data from written documentation and first hand experience


3. Analyse the initiatives

- How do they problematize suicide among Sámi? (what problems are they trying to fix?)
- Use descriptions of initiatives for analysis
- Analyse guided by the "what is the problem represented to be?" (WPR) approach (Bacchi, 2016)



FINDINGS: 17 INITIATIVES

Number of suicide prevention initiatives targeting Sámi in Norway, Sweden, Finland and internationally, per year (2005–2019).



Year

Diversity:


- delivery methods –public meetings and workshops were most common
- target groups
- organizers (but SANKS involved in all Norwegian initiatives)
- levels of adaptation to Sámi context – range from targeting Sámi with not adapted programmes to tailor-made projects

Similarity: few initiatives included thorough evaluation components




UMEÅ UNIVERSITY

WHAT ARE THE "PROBLEMS" IN SAMI SUICIDE PREVENTION INITIATIVES?




IS THE PROBLEM THAT WE DO NOT CHALLENGE NEITHER MAJORITY NOR SÁMI SOCIETIES?





dominant Sámi society

PLAN FOR SUICIDE PREVENTION AMONG SÁMI





"Plan for suicide prevention among Sámi in Norway, Sweden and Finland"


11 strategies:

- Focus on the men
- Produce statistics and support research
- Strengthen Sámi self-determination
- Initiate work to illuminate and deal with historical trauma
- Strengthen and protect Sámi cultural identities
- Decrease violence
- Decrease ethnic discrimination
- Increase diversity and acceptance
- Ensure right to equal health care, adapted to language and culture
- Train and mobilize Sámi civic society
- Strengthen border-crossing cooperation



1: FOCUS ON THE MEN

- Sámi men die more often by suicide than majority men
- Sámi men die by suicide at several times the rate of Sámi women
- Example, Suicide among Sámi in Norway 1970-1998 (Silviken, Haldorsen & Kvernmo 2006):



2: PRODUCE STATISTICS AND SUPPORT RESEARCH

- No official statistics on suicide among Sámi exists because Norway, Sweden and Finland does not register ethnicity
- Available statistics derive from research projects, but does not include all Sámi
- Outdated data:
 - Norway 1970-1998
 - Sweden 1961-2000
 - Finland 1979-2010



3: STRENGTHEN SÁMI SELF-DETERMINATION

- Make sure that Sámi have a real opportunity to influence processes that affects Sámi
- Espescially important for Sámi practicing traditional livelihoods
- Importance stressed by Sámi grassrots




4: INITIATE WORK TO ILLUMINATE AND DEAL WITH HISTORICAL TRAUMA

- Sámi have experienced a wide array of traumatic events throughout history including forced assimilation
- Very little is known about health effects of historical trauma among Sámi
- "Truth and reconcilliation"commissions have started in Norway
 and Finland, as i being planned in
 Sweden


5: STRENGTHEN AND PROTECT SÁMI CULTURAL IDENTITIES

• Ensure that Sámi have access to strong arenas/contexts for Sámi language and culture



6: DECREASE EXPOSURE TO VIOLENCE

• Existing evidence suggest: Sámi at higher risk of exposure to mental, physical and sexual abuse both in childhood and adulthood (Eriksen et al, 2015)



UMEÅ UNIVERSITY

7: DECREASE ETHNIC DISCRIMINATION

- Sámi experience ethnic discrimination at several times the rate of majority populations
- Experiencing discrimination is associated with poor mental health and high suicidality

Suicidality among young Sámi adults in Sweden (Omma et al, 2013)


30%
20%
10%
12,3%
7,3%
3,0%
Suicide plans
Suicide attempt

Discriminated against

UMEÅ UNIVERSITY

8: INCREASE DIVERSITY AND ACCEPTANCE

- Lesbian, gay, bisexual, transgender and queer persons suffers disproportionately from suicidality
- High likelihood of this being true for Sámi LGBTQ-persons as well



9: ENSURE RIGHT TO EQUAL HEALTH CARE, ADAPTED TO LANGUAGE AND CULTURE

- Mental health services are at the core of suicide prevention
- Equal health care demands services adapted to language and cultural needs
- Espescially important for psychiatric services and other "talking cures"
- Availability of such services differs greatly between different Sámi contexts



10: TRAIN AND MOBILIZE SÁMI CIVIC SOCIETY

- "For Sámi by Sámi" holds a partly untapped suicide prevention resource
- This includes training Sámi grass roots but also mobilizing Sámi institutions, organisations etc





11: STRENGTHEN BORDER-CROSSING COOPERATION

- Sámi live in 4 countries, yet suicide preventive efforts often stop by the borders
- When resources are limited, it is more important to pool them



WHERE TO NOW?

- Sweden: new national strategy for mental health and suicide prevention to be developed –
- Ministry for Health and social affairs: Sámi perspectives to be included!
- Collaboration between the Sámi parliament, the Swedish Agency for Public Health, and Department of Epidemiology and Global health, Umeå University:
 - Previous data + rights + strategies
 - o New data:
 - Sámi public health questionnaire (mental health, suicidality)
 - Sámi register study (suicide mortality, patient register)
- Outcome? Hopefully a holistic approach supported by monitoring progress through data!
- Foundation for a Centre for Sámi health at UmU -> research and education







